

Gravitational collapse in asymptotically Anti-de Sitter/de Sitter backgrounds

T. Arun Madhav,^{*} Rituparno Goswami,[†] and Pankaj S. Joshi[‡]

^{*}*Birla Institute of Technology and Science, Pilani*

^{†‡}*Tata Institute of Fundamental Research, Mumbai, India*

We study here the gravitational collapse of a matter cloud with a non-vanishing tangential pressure in the presence of a non-zero cosmological term. Conditions for bounce and singularity formation are derived for the model. It is also shown that when the tangential pressures vanish, the bounce and singularity conditions reduce to that of the dust case studied earlier. The collapsing interior is matched with an exterior which is asymptotically de Sitter or anti de Sitter, depending on the sign of cosmological constant. The junction conditions for matching the cloud to exterior are specified. The effect of the cosmological term on apparent horizons is studied in some detail, and the nature of central singularity is analyzed. We also discuss here the visibility of the singularity and implications for the cosmic censorship conjecture.

PACS numbers: 04.20.Cv, 04.20.Dw, 04.70.Bw

I. INTRODUCTION

The gravitational collapse of a matter cloud which is pressureless dust, and its dynamical evolution as governed by the Einstein's equations, were first studied in detail by Oppenheimer and Snyder [1]. In recent years there have been extensive studies on gravitational collapse to examine the final fate of a collapsing cloud in order to investigate the final state of such a collapse in terms of formation of a black hole or naked singularity. These studies would throw important light on the nature of cosmic censorship and possible mathematical formulations for the same (for some recent reviews, see e.g. [2]). Such studies have already helped to rule out several possible versions of cosmic censorship, where a precise and well-defined formulation itself has been a major unresolved problem so far. While understanding the nature of dynamical gravitational collapse within the framework of Einstein's gravity itself is a problem with considerable astrophysical significance, the understanding of cosmic censorship, if it is valid in some form, is another major motivation for such collapse studies. Most of these investigations so far, however, have assumed a vanishing cosmological term (Λ).

The cosmological constant is sometimes thought of as a constant term in the Lagrangian density of general relativity, and it is also theorized that Λ may be related to the energy density of vacuum (See [3] and references therein). Recent astronomical observations of high red-shift type Ia Supernovae [4] strongly indicate that the universe may be undergoing an accelerated expansion and it is believed that this may be due to a non-vanishing *positive* cosmological constant. On the other hand, the proposed AdS/CFT conjecture [5] in string theory has generated interest in the possibility of space-times with a *negative*

cosmological constant. The conjecture relates string theory in a spacetime where the non-compact part is asymptotically anti-de Sitter (AdS), to a conformal field theory in a space isomorphic to the boundary of AdS. Whereas the obvious effect of a positive cosmological constant is to slow down gravitational collapse, that of the negative cosmological constant is to supplement the gravitational forces.

In this context, it seems pertinent to study the dynamical collapse of matter clouds when $\Lambda \neq 0$. Our purpose here is to study a class of collapsing models which incorporate pressure, and which are asymptotically either de Sitter or anti-de Sitter geometry, depending on the sign of the cosmological term. Dust models with Λ are known in literature [6] and there have been some studies on dust collapse with Λ in recent years (see e.g. [7]). We intend to discuss here a sufficiently general fluid model which allows pressure to be non-zero, and which also allows the cosmological term to be non-vanishing. Specifically, we study models with a non-vanishing *tangential pressure* p_θ [8], together with $\Lambda \neq 0$. Collapse models with a tangential pressure have been studied extensively, but not with a non-zero Λ [9]. Allowing the collapse to develop from regular initial conditions, we study the bounce and singularity formation conditions, junction conditions at the boundary of the cloud so as to match it to a suitable exterior, and we also consider how the apparent horizons are affected by the presence of Λ . One is also interested in considering the possibility whether the presence of Λ could restore the *cosmic censorship conjecture* (see e.g. [10] and references therein), and to investigate how the final end state of gravitational collapse is affected by Λ . The present model generalizes the Lemaître-Tolman-Bondi (LTB) dust [11] model studies with Λ , by introducing non-zero pressures in the collapsing cloud.

The relevant form of Einstein equations, conditions to ensure that the collapse develops from a regular initial data, and the necessary energy conditions are introduced in Section II, together with the details of the tangential pressure model. In Section III, we derive the evolution of the collapsing matter shells, and explicitly give the

^{*}Electronic address: arunmadhav@gmail.com

[†]Electronic address: goswami@tifr.res.in

[‡]Electronic address: psj@tifr.res.in

conditions when a singularity is formed and when the bounce of a particular shell occurs during the collapse evolution. The reduction to $\Lambda \neq 0$ dust collapse case, when the tangential pressures are put to zero, is also demonstrated. For the collapsing solution to be physically plausible, it must satisfy certain junction conditions at the boundary hypersurface where the interior collapsing cloud joins with a suitable exterior spacetime. In Section IV we study the matching of the collapsing interior to the exterior Schwarzschild de Sitter or anti-de Sitter spacetime, and in Section V we discuss in brief the effect of Λ on the apparent horizons of the fluid model. In section VI the nature of the singularity in the tangential pressure fluid and dust models is considered when $\Lambda \neq 0$, in terms of its being hidden within a black hole, or whether it would be visible to outside observers. The final Section VII summarizes some conclusions.

II. EINSTEIN EQUATIONS, REGULARITY AND ENERGY CONDITIONS

The general spherically symmetric metric in the co-moving coordinates (t, r, θ, ϕ) is given as,

$$ds^2 = -e^{2\nu(t,r)} dt^2 + e^{2\psi(t,r)} dr^2 + R^2(t,r) d\Omega^2 \quad (1)$$

where $d\Omega^2 = d\theta^2 + \sin^2\theta d\phi^2$ is the line element on a two-sphere. The metric variables $e^{2\nu(t,r)}$ and $e^{2\psi(t,r)}$ are functions of t and r . The energy-momentum tensor for a general *Type I* matter field [12] can be written as,

$$T_t^t = -\rho; T_r^r = p_r; T_\theta^\theta = T_\phi^\phi = p_\theta \quad (2)$$

The quantities ρ , p_r and p_θ are the matter density, radial pressure and the tangential pressure respectively. It is assumed that the matter field satisfies the *weak energy condition* [12]. It means that the energy density as measured by any local observer is positive. Then, for any timelike vector V^μ we have,

$$T_{\mu\gamma} V^\mu V^\gamma \geq 0 \quad (3)$$

This amounts to,

$$\rho + \Lambda \geq 0; \rho + p_r \geq 0; \rho + p_\theta \geq 0 \quad (4)$$

The evolution of the matter cloud is determined by the Einstein equations and for the metric (1) these are given by (in the units of $8\pi G = c = 1$),

$$\rho + \Lambda = \frac{F'}{R^2 R'}, \quad p_r - \Lambda = -\frac{\dot{F}}{R^2 \dot{R}} \quad (5)$$

$$\nu'(\rho + p_r) = 2(p_\theta - p_r) \frac{R'}{R} - p_r' \quad (6)$$

$$-2\dot{R}' + R' \frac{\dot{G}}{G} + \dot{R} \frac{H'}{H} = 0 \quad (7)$$

$$G - H = 1 - \frac{F}{R} \quad (8)$$

where $(\dot{})$ and $()'$ represent partial derivatives with respect to t and r respectively and,

$$G(r, t) = e^{-2\psi(R')^2}, \quad H(r, t) = e^{-2\nu} \dot{R}^2 \quad (9)$$

Here $F(r, t)$ is an arbitrary function, and in spherically symmetric spacetimes, it has the interpretation of the *mass function* of the collapsing cloud, in the sense that this represents the total gravitational mass within a shell of comoving radius r [13]. The boundary of the collapsing cloud is labelled by the comoving coordinate r_π . In order to preserve regularity at initial epoch we require $F(t_i, 0) = 0$.

It can be seen from equation (5) that the density of the matter blows up when $R = 0$ or $R' = 0$. The case $R' = 0$ corresponds to *shell-crossing* singularities. The shell-cross singularities are generally considered to be weak and possibly removable singularities. Hence we shall consider here only the *shell-focusing* singularities (taking $R' > 0$), where the physical radius of all the matter shells go to a zero value ($R = 0$). Let us use the scaling independence of the coordinate r to write,

$$R(t, r) = rv(t, r) \quad (10)$$

where ' v ' is the *Scale factor*. We have,

$$v(t_i, r) = 1; \quad v(t_s(r), r) = 0; \quad \dot{v} < 0 \quad (11)$$

where t_i and t_s stand for the initial and the singular epochs respectively. This means we scale the radial coordinate r in such a way that at the initial epoch $R = r$, and at the singularity, $R = 0$. The condition $\dot{v} < 0$ signifies that we are dealing with gravitational collapse. From the point of view of initial data, at the initial epoch $t = t_i$, we have five functions of coordinate r given by $\nu_0(r)$, $\psi_0(r)$, $\rho_0(r)$, $p_{r_0}(r)$ & $p_{\theta_0}(r)$. Note that initial data are not all mutually independent. To preserve regularity and smoothness of initial data we must make some assumptions about the initial pressures at the regular center $r = 0$. Let the gradients of pressures vanish at the center, that is, $p_{r_0}'(0) = p_{\theta_0}'(0) = 0$. The difference between radial and tangential pressures at the center should also vanish, i.e. $p_{r_0}(0) - p_{\theta_0}(0) = 0$. It is seen that we have a total of five field equations with seven unknowns, ρ , p_r , p_θ , ψ , ν , R , and F , giving us the freedom to choose two free functions. Selection of these functions, subject to the given initial data and weak energy condition, determines the matter distribution and metric of the space-time and thus leads to a particular dynamical evolution of the initial data.

Spherically symmetric collapse models, where the radial pressure is taken to be vanishing, but the tangential pressure could be non-zero have been studied in some detail over the past few years [9]. The main motivation in

the present consideration is to study bounce and singularity formation conditions for the case when we have a non-vanishing Λ term present, when pressures are also introduced and allowed to be non-zero within a collapsing cloud. One would also like to understand how Λ affects the junction conditions at the boundary where the cloud is matched to an exterior spacetime, the horizon formation, and the nature of central singularity, when non-zero pressures are present within the cloud.

Taking $p_r = 0$ in Eq. (5) gives $F(t, r)$ and ρ in the following forms,

$$F(t, r) = r^3 \mathcal{M}(r) + \frac{\Lambda}{3} R^3 \quad (12)$$

$$\rho = \frac{3\mathcal{M} + r [\mathcal{M}_{,r}]}{v^2(v + rv')} \quad (13)$$

Here $\mathcal{M}(r)$ is an arbitrary function of r subject to the energy conditions. There remains the freedom to choose one function, since there are six equations with seven unknowns. In order to work within the framework of a specific class of models, we take $\nu(t, r)$ in the specific form,

$$\nu(t, r) = c(t) + \nu_0(R) \quad (14)$$

The conditions imposed here, namely that of vanishing radial pressures, and Eq. (14) specifying a form of ν may be considered to be strong assumptions. However, this enables us to make our study in sufficient generality with sufficiently rich structure as we shall see below, with non-zero pressures introduced into the collapse model. A mechanism by which we can have non-vanishing tangential pressures is illustrated by the Einstein cluster [14]. This is a spherically symmetric cluster of rotating particles where the motion of the particles is sustained by an angular momentum. This has an average effect and creates non-zero tangential stresses within the collapsing cloud.

Also, one can rederive the dust collapse models with a non-vanishing cosmological term, by putting $\nu_0(R) = 0$ in Eq. (14) and redefining the comoving time coordinate. It is thus clear that the class of models considered generalize the dust collapse models with a non-vanishing cosmological constant. In general, as $v \rightarrow 0$, $\rho \rightarrow \infty$. Thus the density blows up at the singularity $R = 0$ which will be a curvature singularity as expected. Using Eq. (14) in Eq. (7), we have,

$$G(t, r) = b(r)e^{2\nu_0(R)} \quad (15)$$

Here $b(r)$ is an arbitrary, C^2 function of r . In correspondence with the dust models, we can write,

$$b(r) = 1 + r^2 b_0(r) \quad (16)$$

where $r^2 b_0(r)$ is the *energy distribution function* for the collapsing shells. Using the given initial data and Eq. (6)

one can obtain the function $\nu_0(R)$. Substituting Eq. (14) in Eq. (6) we get the intrinsic equation of state,

$$p_\theta = \frac{R}{2} \nu_{,R} \rho \quad (17)$$

Finally, using equations (12), (14) and (15) in Eq. (8), we have,

$$\sqrt{R} \dot{R} = -a(t)e^{\nu_0(R)} \sqrt{(1 + r^2 b_0) R e^{2\nu_0} - R + r^3 \mathcal{M} + \frac{\Lambda}{3} R^3} \quad (18)$$

Here $a(t)$ is a function of time and by a suitable scaling of the time coordinate, we can always make $a(t) = 1$. We deal here with the collapse models, and so the negative sign is due to the fact that $\dot{R} < 0$, which represents the collapse condition.

III. SINGULARITY FORMATION AND REBOUNCE

One of our main purposes here is to examine how the introduction of a non-vanishing cosmological term modifies the collapse dynamics. For example, in the case of dust collapse, once the collapse initiates from an initial epoch, there cannot be any reversal or a bounce, and the gravity forces the cloud to collapse necessarily to a singularity. However, this need not be the case when the cosmological term is non-vanishing, and we have to re-examine the collapse dynamics in order to find how the collapse evolves. This we do here for the particular class of tangential collapse models as specified above, which also generalizes the dust collapse case.

The evolution of a particular shell may be deduced from Eq. (18). Rewriting Eq. (18) in terms of the scale factor we have,

$$\dot{v}^2 = \frac{e^{2\nu_0} [v j(r, v) + \mathcal{M} + \frac{\Lambda}{3} v^3]}{v} = V(r, v) \quad (19)$$

where

$$j(r, v) = \frac{b(r)e^{2\nu_0(rv)} - 1}{r^2} \quad (20)$$

The right hand side of the equation (19) may be thought of as an *effective potential* ($V(r, v)$) for a shell. The allowed regions of motion correspond to $V(r, v) \geq 0$, as $\dot{v}^2$ is non-negative, and the dynamics of the shell may be studied by finding the turning points. If we start from an initially collapsing state ($\dot{v} < 0$), we will have a rebound if we get $\dot{v} = 0$, before the shell has become singular. This can happen when $V(r, v) = 0$. Hence, to study the various evolutions for a particular shell we must analyze the roots of the equation $V(r, v) = 0$ keeping the value r to be fixed. It will be seen that the cosmological constant appearing in the effective potential does play an important role in the evolution of a shell.

To clarify these ideas, let us consider a smooth initial data, where the initial density, pressure, and energy distributions are expressed as only even powers of r . Such a consideration, that the initial data be smooth, is often justified on physical grounds. So we take,

$$\rho(t_i, r) = \rho_{00} + \rho_2 r^2 + \rho_4 r^4 + \dots \quad (21)$$

$$p_\theta(t_i, r) = p_{\theta_2} r^2 + p_{\theta_4} r^4 + \dots \quad (22)$$

$$b_0(r) = b_{00} + b_{02} r^2 + \dots \quad (23)$$

With the above form of smooth initial data to evolve in time using the Einstein equations, we can explicitly integrate Eq. (6) at the initial epoch to get,

$$\nu_0(R) = p_{\theta_2} R^2 + \frac{(p_{\theta_4} - \rho_2 p_{\theta_2})}{2} R^4 + \dots \quad (24)$$

We have neglected here higher order terms in the expansion, since at present we want to concentrate on the evolution of shells near $r = 0$. The conditions when the treatment is applicable to the whole cloud will be discussed later. Eq. (19) near the center of the cloud ($r \ll r_\pi$) may be written as,

$$\dot{v}^2 = \frac{(1 + 2p_{\theta_2} r^2 v^2) [(\Lambda + 6p_{\theta_2} b(r)) v^3 + 3b_0(r) v + 3\mathcal{M}]}{3v} \quad (25)$$

The first factor in $V(r, v)$ is initially positive, because it is the $|g_{00}|$ term. As the collapse evolves, the scale factor (v) reduces from 1 at the initial epoch to 0 at the time of singularity. Hence it clear that the first factor can never become zero, and hence does not contribute to a bounce of the shell. The main features of the evolution of the cloud basically derive from the second factor in $V(r, v)$.

The second factor in the effective potential expression is a cubic equation which in general has three roots. Only positive real roots correspond to physical cases. Since the coefficient in the second power is zero, we may conclude that if all three roots are real then at least one of them has to be positive and at least one negative. We observe that $V(r, 0) = \mathcal{M} > 0$. Hence, any region between $R = rv = 0$ and the first positive zero of $V(r, v)$ always becomes singular during collapse. The region between the unique positive roots is forbidden since in those regions $\dot{v}^2 < 0$. For a particular shell to bounce it must therefore lie, during initial epoch ($v = 1$), in a region to the right of the second positive root. We will now analyze the various cases for $\Lambda \neq 0$ in detail and derive the necessary conditions.

1. If

$$b_0(r) \geq 0 ; (\Lambda + 6p_{\theta_2} b(r)) > 0 \quad (26)$$

then from the Descartes's rule of signs (See for example [15]) we see that there are no positive roots. Thus a *singularity always forms* from initial collapse.

2. If

$$b_0(r) \geq 0 ; (\Lambda + 6p_{\theta_2} b(r)) < 0 \quad (27)$$

we infer from the sign rule that there is exactly one positive root ($\alpha(r)$). The other two roots are negative or complex conjugates. The allowed space of dynamics is $[0, \alpha]$. Thus $\alpha \geq 1$ would always ensure a singularity. However $\alpha < 1$ implies an unphysical situation initially, where $\dot{v}^2 < 0$.

3. If

$$b_0(r) \leq 0 ; (\Lambda + 6p_{\theta_2} b(r)) < 0 \quad (28)$$

there is exactly one positive root ($\beta(r)$). Again all shells in the allowed dynamical space $[0, \beta]$ become *singular* starting from initial collapse.

4. If

$$b_0(r) < 0 ; (\Lambda + 6p_{\theta_2} b(r)) > 0 \quad (29)$$

there are three possibilities,

4.1 If

$$\mathcal{M}^2 > -4 \frac{b_0^3(r)}{9(\Lambda + 6p_{\theta_2} b(r))} \quad (30)$$

then there are no positive roots and a *singularity* is always the final outcome of collapse for shells under consideration.

4.2 If

$$\mathcal{M}^2 < -4 \frac{b_0^3(r)}{9(\Lambda + 6p_{\theta_2} b(r))} \quad (31)$$

then there are two positive roots ($\gamma_1(r)$ and $\gamma_2(r)$ respectively). The space of allowed dynamics is $[0, \gamma_1]$ and $[\gamma_2, \infty)$. The region (γ_1, γ_2) is forbidden. Shells in the $[0, \gamma_1]$ region initially, *always become singular*. But shells initially belonging to the region $[\gamma_2, \infty)$ will *undergo a bounce and subsequent expansion* starting from initial collapse. This bounce occurs when their geometric radius approaches $R_{\text{bounce}} = r\gamma_2$. Using the definitions

$$\varrho = \sqrt{\frac{-4b_0(r)}{\Lambda + 6p_{\theta_2} b(r)}} \quad (32)$$

$$\vartheta = \frac{1}{3} \text{Cos}^{-1} \left[-\sqrt{\frac{-9\mathcal{M}(r)^2(\Lambda + 6p_{\theta_2} b(r))}{4b_0^3(r)}} \right] \quad (33)$$

the condition for a particular shell to become singular or undergo a bounce may be explicitly written in terms of the initial data and Λ as,

$$1 < \varrho \text{Cos}\vartheta ; \text{Singularity} \quad (34)$$

$$1 > \varrho \text{Cos}(\vartheta + \frac{4\pi}{3}) ; \text{Bounce} \quad (35)$$




FIG. 1: The effective potential profile ($V(r, v)$) for the $r = 0$ shell with $(\Lambda + 6p_{\theta_2}) = 7.1978 \times 10^{-3}$, $b_{00} = -3 \times 10^{-3}$, and $\mathcal{M} = 1.033 \times 10^{-3}$ in appropriate units.

Here $\varrho \cos \vartheta$ is $\gamma_1(r)$ and $\varrho \cos(\vartheta + \frac{4\pi}{3})$ is $\gamma_2(r)$, which are the two roots of the potential function. Note that contrary to the dust models with Λ , there may be a bounce for both positive and negative values of the cosmological constant.

4.3 If

$$\mathcal{M}^2 = -4 \frac{b_0^3(r)}{9(\Lambda + 6p_{\theta_2}b(r))} \quad (36)$$

the positive roots are equal. There is no forbidden region, and there will be a bounce if $1 > \frac{\varrho}{2}$. Fig. 1. illustrates a particular choice of initial data and Λ which causes a bounce in the central ($r = 0$) shell.

To analyze the evolution of shells far from the center, in general one has to resort to numerical methods, and it is difficult to analytically give a simple expression for the singularity or bounce conditions. Nevertheless, the analysis as given here becomes valid for the entire cloud all the way till boundary when the geometrical radius of the cloud boundary at the initial epoch ($R(t_i, r_\pi) = r_\pi$) is itself small relative to the initial data coefficients (ie. $\rho_n r_\pi^n, p_{\theta_n} r_\pi^n \ll 1$). Also, if we choose the initial data such that the higher coefficients in the power series expansion are zero (i.e. $p_{\theta_n \geq 4}, \rho_{k \geq 4} = 0$) and $\rho_2 r_\pi^2, p_{\theta_2} r_\pi^2 \ll 1$, then the analysis is again applicable to the whole cloud. Thus the results derived can be considered quite general in these circumstances and applicable to the cloud as a whole.

In this context, if all shells in the collapsing cloud satisfy Eq. (29), (31) and (35), the complete cloud undergoes bounce starting from initial collapse. To avoid shell crossings the sufficient condition would be,

$$\forall r \in [0, r_\pi) \quad , \quad \gamma_2(r + \delta) \geq \gamma_2(r) \quad (37)$$

where δ is an infinitesimal increment in the comoving radius. It is seen that in all the cases discussed, it is not Λ alone, but $\Lambda + 6p_{\theta_2}b(r)$ along with $b_0(r)$ that determines the evolution of the shell. This is in contrast to

the dust models with a non-zero Λ , where solely the cosmological constant decided the evolution of a shell for a given energy function. It is also interesting to note that unlike the dust collapse models with Λ , there could be a bounce in the fluid model with vanishing radial pressures for both positive and negative values of the cosmological constant. This is due to the contribution from the tangential pressure.

It can be seen now that one can rederive the known bounce conditions in $\Lambda \neq 0$ dust collapse case (Deshingkar *et al* [7]) as a special case of the consideration here, when $p_\theta = 0$. In that case,

$$\dot{v}^2 = \frac{\Lambda v^3 + 3b_0(r)v + 3\mathcal{M}}{3v} = V(r, v) \quad (38)$$

The study now becomes valid for all shells, without any approximation. For example, following similar steps as for the fluid model, one obtains for $\Lambda > 0$,

1. If

$$b_0(r) > 0 \quad (39)$$

the singularity always forms from initial collapse.

2. If

$$b_0(r) = 0 \quad (40)$$

then again it is found that all shells become singular from collapse.

3. If

$$b_0(r) < 0 \quad (41)$$

there are two scenarios possible. For

$$\mathcal{M}^2 > -4 \frac{b_0^3}{9\Lambda} \quad (42)$$

all shells become singular. For

$$\mathcal{M}^2 < -4 \frac{b_0^3}{9\Lambda} \quad (43)$$

there are two positive roots (ψ_1 & ψ_2 respectively) for $V(r, v)$. Shells belonging to $[0, \psi_1]$ always become singular while those belonging to $[\psi_2, \infty)$ undergo bounce starting from initial collapse.

IV. SPACETIME MATCHING

As we pointed out above, there is a strong physical motivation to study and investigate the gravitational collapse phenomena in a background which is asymptotically either a de Sitter or anti-de Sitter metric. For this purpose, the collapsing cloud has to be matched at the

boundary to a suitable exterior spacetime which has the desired properties.

In the present case, we shall show below that the exterior vacuum spacetime of the collapsing region may be described by the Schwarzschild- de Sitter (SdS) or Schwarzschild- Anti de Sitter (SAdS) metric, depending on whether the cosmological constant is taken to be positive or negative. The collapsing interior cloud which has a non-zero tangential pressure is then to be smoothly matched to an exterior spacetime in order to generate the full spacetime. The necessary and sufficient conditions to achieve a smooth matching are given by the Israel-Darmois junction conditions ([16],[17]), which we shall use below.

Let the interior of the collapsing cloud be described by the metric,

$$S^- : ds_-^2 = -e^{2\nu(t,r)} dt^2 + e^{2\psi(t,r)} dr^2 + R^2(t,r) d\Omega^2 \quad (44)$$

The exterior vacuum solution can be given as the Schwarzschild-de Sitter/Anti de Sitter spacetime (as decided by the sign chosen for the cosmological term) as given by,

$$S^+ : ds_+^2 = -\mathcal{D}(\mathcal{R}) dT^2 + \mathcal{D}^{-1}(\mathcal{R}) d\mathcal{R}^2 + \mathcal{R}^2 d\Omega^2 \quad (45)$$

where $\mathcal{D}(\mathcal{R})$ is given by,

$$\mathcal{D}(\mathcal{R}) = 1 - \frac{2M}{\mathcal{R}} \mp \frac{|\Lambda|\mathcal{R}^2}{3} \quad (46)$$

The negative sign precedes $|\Lambda|$ for a SdS exterior and the positive for SAdS exterior. Note that as $\mathcal{R} \rightarrow \infty$, the $\mp \frac{|\Lambda|\mathcal{R}^2}{3}$ term dominates over the $\frac{2M}{\mathcal{R}}$ term, and the spacetime approaches asymptotically de Sitter metric ($-\frac{|\Lambda|\mathcal{R}^2}{3}$), or the Anti-de Sitter metric ($+\frac{|\Lambda|\mathcal{R}^2}{3}$), as the case may be.

Let Π denote the boundary hypersurface. The equations of the boundary hypersurface considered as an embedding in the interior or exterior spacetimes are,

$$\Pi^- : r - r_\pi = 0 \quad \Pi^+ : \mathcal{R} - \mathcal{R}_\pi(T) = 0 \quad (47)$$

Substituting (47) in (44) and (45) we get the metric on the hypersurface as,

$$S_\Pi^- : ds_-^2 = -e^{2\nu(t,r_\pi)} dt^2 + R^2(t,r_\pi) d\Omega^2 \quad (48)$$

$$S_\Pi^+ : ds_+^2 = -\mathcal{D}(\mathcal{R}_\pi) dT^2 + \mathcal{D}^{-1}(\mathcal{R}_\pi) d\mathcal{R}_\pi^2 + \mathcal{R}_\pi^2 d\Omega^2 \quad (49)$$

The Israel-Darmois conditions to match the interior spacetime with exterior require that the first and second fundamental forms of the boundary hypersurface match. The first fundamental form is given by,

$$g_{\mu\nu} d\zeta^\mu d\zeta^\nu \quad (50)$$

where ζ parametrizes the hypersurface ($\zeta^t : \tau, \theta, \phi$). The matching of the first fundamental form gives, from Eq. (48) and Eq. (49),

$$R(t, r_\pi) = \mathcal{R}_\pi \quad (51)$$

$$d\tau = e^{\nu(t,r_\pi)} dt \quad (52)$$

$$d\tau = \left[\mathcal{D}_\pi - \frac{\mathcal{R}_{\pi,T}^2}{\mathcal{D}_\pi} \right]^{1/2} dT \quad (53)$$

The above three conditions must be satisfied for a smooth matching of the collapsing interior to the exterior spacetime. The next set of conditions will be given by the matching of the second fundamental forms.

The second fundamental form is given by

$$K_{\mu\nu} d\zeta^\mu d\zeta^\nu \quad (54)$$

where,

$$K_{\mu\nu}^\pm = -n_\sigma^\pm x_{,\zeta^\mu, \zeta^\nu}^\sigma - n_\sigma^\pm \Gamma_{\beta\gamma}^\sigma x_{,\zeta^\mu}^\beta x_{,\zeta^\nu}^\gamma \quad (55)$$

is the extrinsic curvature [18]. The $(,)$ denotes partial differentiation. n_σ is the normal to the hypersurface which is given by,

$$n_\sigma = \frac{f_{,\sigma}}{[g^{\mu\nu} f_{,\mu} f_{,\nu}]^{1/2}} \quad (56)$$

where $f = 0$ is the equation of the boundary hypersurface. Direct calculation gives,

$$n_\sigma^- = (0, e^{\psi(t,r_\pi)}, 0, 0) \quad (57)$$

$$n_\sigma^+ = (-\mathcal{R}_{,\tau}^\pi, T_{,\tau}, 0, 0) \quad (58)$$

The $K_{\theta\theta}$ extrinsic curvatures are calculated as,

$$K_{\theta\theta}^+ = (\mathcal{D}\mathcal{R}T_{,\tau})_\pi \quad K_{\theta\theta}^- = (RR_{,r}e^{-\psi})_\pi \quad (59)$$

Now using the fact that the second fundamental forms match (i.e. $[K_{\theta\theta}^+ - K_{\theta\theta}^-]_\pi = 0$), we get using equations (51),(52),(53) and (59) after simplification,

$$R_{\pi,r}^2 e^{-2\psi} - R_{\pi,\tau}^2 e^{-2\nu} = 1 - \frac{2M}{\mathcal{R}_\pi} \mp \frac{|\Lambda|}{3} \mathcal{R}_\pi^2 \quad (60)$$

This is identical to the Cahill and McVittie definitions for the mass function ([13],[19]), and hence from Eq. (8) we may by comparison take

$$F_\pi = 2M \pm \frac{|\Lambda|}{3} \mathcal{R}_\pi^3 \quad (61)$$

This expression suggests that for a smooth matching of the interior and exterior spacetimes the interior mass function at the surface must equal the generalized Schwarzschild mass.

As we can see there is a contribution to the mass function from the cosmological constant. A positive cosmological constant has an additive contribution and a negative cosmological constant has a deductive contribution to the mass function F_π . The matching of a collapsing dust interior with exterior SdS/SAdS would give similar results [7]). The $K_{\tau\tau}$ and $K_{\phi\phi}$ components of the extrinsic curvature may similarly be calculated. The condition $[K_{\tau\tau}^+ - K_{\tau\tau}^-]_\pi = 0$ gives no new information since the radial pressure is zero. Due to spherical symmetry, $[K_{\phi\phi}^+ - K_{\phi\phi}^-]_\pi = 0$ gives the same result as Eq. (61). All the above conditions (51), (52), (53) and (61) must be satisfied for smooth matching of spacetimes across the boundary.

V. HORIZONS

Apparent horizons ($\mathcal{H}$) are the boundaries of trapped regions [12] in the spacetime. We discuss below the effect of a non-zero Λ term on the apparent horizons of the tangential pressure fluid collapse models considered here briefly. In general, the equation of $\mathcal{H}$ can be written as,

$$\mathcal{H} : g^{\mu\nu} R_{,\mu} R_{,\nu} = 0 \quad (62)$$

Substituting (44) in (62) we get,

$$R_{,r}^2 e^{-2\psi} - R_{,\tau}^2 e^{-2\nu} = 0 \quad (63)$$

From the definition of the mass function Eq. (8), and (63) we therefore have,

$$1 - \frac{F}{R} = 0 \quad (64)$$

Finally, from (12) and (64),

$$\mathcal{H} : (3 - \Lambda R^2) R = 3r^3 \mathcal{M} \quad (65)$$

When $\Lambda = 0$ there is necessarily only one apparent horizon given by

$$R = r^3 \mathcal{M} \quad (66)$$

which is the Schwarzschild horizon in case we are considering that geometry. The same equation also defines horizon within the collapsing cloud, where $R(t, r)$ is one of the metric functions. For the case when $\Lambda > 0$, Eq. (65) is a cubic equation with at least one positive and one negative root. For $\Lambda > 0$ the various cases are given as below.

1. For

$$3r^3 \mathcal{M} < \frac{2}{\sqrt{\Lambda}} \quad (67)$$

there are two positive roots for (65) and hence there are *two* apparent horizons. These horizons are given by

$$R_c(r) = \frac{2}{\sqrt{\Lambda}} \cos \left[\frac{1}{3} \cos^{-1} \left(-\frac{3}{2} r^3 \mathcal{M} \sqrt{\Lambda} \right) \right] \quad (68)$$

$$R_b(r) = \frac{2}{\sqrt{\Lambda}} \cos \left[\frac{4\pi}{3} + \frac{1}{3} \cos^{-1} \left(-\frac{3}{2} r^3 \mathcal{M} \sqrt{\Lambda} \right) \right] \quad (69)$$

These have been at times called the cosmological, and the black hole horizons [20].

2. For

$$3r^3 \mathcal{M} = \frac{2}{\sqrt{\Lambda}} \quad (70)$$

there is only one positive root for (65), given by

$$R_{bc}(r) = \frac{1}{\sqrt{\Lambda}} \quad (71)$$

This corresponds to a *single apparent horizon*.

3. For

$$3r^3 \mathcal{M} > \frac{2}{\sqrt{\Lambda}} \quad (72)$$

there are no positive roots and hence there are *no apparent horizons*.

The case (72) also shows that the mass of the black hole is bounded above by $F = \frac{1}{\sqrt{\Lambda}}$ and attains the largest proper area $4\pi/\Lambda$. A general result exists in literature [20] showing that in spacetimes with $\Lambda > 0$ and matter satisfying the strong energy condition, the area of a black hole cannot exceed $4\pi/\Lambda$. A detailed treatment of apparent horizons in the Lemaitre-Tolman-Bondi dust collapse models with a non-zero Λ term is given by Cissoko M. *et al* [7].

From Eq. (18), the time of apparent horizon formation in the fluid model is given by

$$t_{ah}(r) = t_s(r) - t_1(r) \quad (73)$$

where $t_1(r)$ is defined as,

$$t_1(r) = \int_0^{v_{ah}(r)} \frac{\sqrt{v} dv}{\sqrt{e^{4\nu_0} v b_0 + e^{2\nu_0} (v^3 h(rv) + \mathcal{M} + \frac{\Lambda}{3} v^3)}} \quad (74)$$

Also $rv_{ah} = R_b$ or R_c , $h(R) = (e^{2\nu_0(R)} - 1)/R^2$ and $t_s(r)$ is the time of singularity formation. For shells close to the center of the cloud ($r \ll r_\pi$) the expression becomes,

$$t_{ah}(r) = t_s(r) - \int_0^{v_{ah}(r)} \frac{\sqrt{3v} dv}{\sqrt{(\Lambda + 6p_{\theta_2}) v^3 + 3b_0(r) v + 3\mathcal{M}}} \quad (75)$$

It is observed that the cosmological constant modifies the time of *formation of horizons* and also the *time lag* between horizon formation and singularity formation in the fluid model.

VI. NATURE OF THE CENTRAL SINGULARITY

The final end state of gravitational collapse and the nature of the resulting singularity continue to be among the most outstanding problems in gravitation theory and relativistic astrophysics today. As pointed out earlier, the hypothesis that such a collapse leading to a singularity, under physically realistic conditions must end in the formation of a black hole, and that the eventual singularity must be hidden below the event horizons of gravity is the *cosmic censorship conjecture*. Despite numerous attempts, this conjecture as such remains a major unsolved problem lying at the foundation of black hole physics today.

From such a perspective, we need to examine the nature of the singularity, in terms of its visibility or otherwise for outside observers, when it develops within the context of the models considered here. This should tell us how the presence of the Λ term modifies these considerations, because we already know that in the case of a tangential pressure present, but a vanishing cosmological constant, both black holes and naked singularities do develop as final collapse end states depending on the nature of the initial data (see e.g. [8]).

We have already seen above that the cosmological constant modifies the time of formation of trapped surfaces. If the formation of the horizon precedes the formation of the central singularity then the singularity will be necessarily covered, i.e. it is a black hole. If on the other hand, the horizon formation occurs after the singularity formation, there may be future directed non-spacelike geodesics that end in the past at the singularity. Then the final end state would be a naked singularity. Thus we need to find whether there exist future directed null geodesics that end at the singularity in the past.

Towards analysing this issue, let us define a function $h(R)$ as,

$$h(R) = \frac{e^{2\nu_0(R)} - 1}{R^2} = 2g(R) + \mathcal{O}(R^2) \quad (76)$$

Using Eq. (76) in Eq. (18), we get after simplification,

$$\sqrt{v}\dot{v} = -\sqrt{e^{4\nu_0}vb_0 + e^{2\nu_0}\left(\left(\frac{\Lambda}{3} + h(rv)\right)v^3 + \mathcal{M}\right)} \quad (77)$$

Integrating the above equation, we get,

$$t(v, r) = \int_v^1 \frac{\sqrt{v}dv}{\sqrt{e^{4\nu_0}vb_0 + e^{2\nu_0}\left(v^3\left(h(rv) + \frac{\Lambda}{3}\right) + \mathcal{M}\right)}} \quad (78)$$

The time of formation of a shell focussing singularity, for a specific shell, is obtained by taking the limits of integration in above as (0,1). The shells collapse consecutively, one after the other to the center as there are no shell-crossings ($R' > 0$). We are interested in the central shell (i.e. the singularity forming at $r = 0$), since we will see

that all $r > 0$ shells are necessarily covered on becoming singular. Taylor expanding the above function around $r = 0$, we get,

$$t(v, r) = t(v, 0) + r \left. \frac{dt(v, r)}{dr} \right|_{r=0} + \frac{r^2}{2!} \left. \frac{d^2t(v, r)}{d^2r^2} \right|_{r=0} + \dots \quad (79)$$

Let us denote,

$$\mathcal{X}_n(v) = \left. \frac{d^nt(v, r)}{dr^n} \right|_{r=0} \quad (80)$$

The initial data is taken to be smooth (ie. with only even powers of r allowed). Due to this choice, the first derivative of the functions appearing in the above equation vanish at $r = 0$. Hence we have,

$$\mathcal{X}_1(v) = 0 \quad (81)$$

Defining $B_f(r, v) = e^{4\nu_0}vb_0 + e^{2\nu_0}\left(\left(\frac{\Lambda}{3} + h\right)v^3 + \mathcal{M}\right)$ we may write the Taylor expansion about $r = 0$ as,

$$t(v, r) = t(v, 0) - \frac{r^2}{4} \int_v^1 \frac{B_f''(0, v)\sqrt{v}dv}{B_f(0, v)^{3/2}} + \dots \quad (82)$$

and we have,

$$\mathcal{X}_2(v) = - \int_v^1 \frac{B_f''(0, v)\sqrt{v}dv}{B_f(0, v)^{3/2}} \quad (83)$$

In order to consider the possibility of existence of null geodesic families which end at the singularity in the past, and to examine the nature of the singularity occurring at $R = 0, r = 0$ in this model, let us consider the outgoing null geodesic equation which is given by,

$$\frac{dt}{dr} = e^{\psi-\nu} \quad (84)$$

We use a method which is similar to that given in [21]. The singularity curve is given by $v(t_s(r), r) = 0$, which corresponds to $R(t_s(r), r) = 0$. Therefore, if we have any future directed outgoing null geodesics terminating in the past at the singularity, we must have $R \rightarrow 0$ as $t \rightarrow t_s$ along the same. Now writing Eq. (84) explicitly in terms of variables ($u = r^\alpha, R$), we have,

$$\frac{dR}{du} = \frac{1}{\alpha} r^{-(\alpha-1)} R' \left[1 - \sqrt{\frac{be^{2\nu_0} + \frac{r^3\mathcal{M}}{R} + \frac{\Lambda R^2}{3} - 1}{be^{2\nu_0}}} \right] \quad (85)$$

Eq. (85) is required to be *finite and positive*, for the existence of a naked singularity [21]. In order to get the tangent to the null geodesic in the (R, u) plane, we choose a particular value of α such that the geodesic equation is expressed only in terms of $\left(\frac{R}{u}\right)$. A specific value of α is to be chosen which enables us to calculate the proper limits at the central singularity. In the tangential pressure collapse model discussed in the previous section we have $\mathcal{X}_1(0) = 0$, and hence we choose $\alpha = \frac{2}{3}$ so that when the

limit $r \rightarrow 0, t \rightarrow t_s$ is taken we get the value of tangent to null geodesic in the (R, u) plane as,

$$\frac{dR}{du} = \frac{3}{7} \left(\frac{R}{u} + \frac{\sqrt{M_0} \mathcal{X}_2(0)}{\sqrt{\frac{R}{u}}} \right) \frac{(1 - \frac{F}{R})}{\sqrt{G}(\sqrt{G} + \sqrt{H})} \quad (86)$$

Now note that for any point with $r > 0$ on the singularity curve $t_s(r)$ we have $R \rightarrow 0$ whereas, F (interpreted as the gravitational mass within the comoving radius r) tends to a finite positive value once the energy conditions are satisfied. Under the situation, the term F/R diverges in the above equation, and all such points on the singularity curve will be covered as there will be no outgoing null geodesics from such points.

We hence need to examine the central singularity at $r = 0, R = 0$ to determine if it is visible or not. That is, we need to determine if there are any solutions existing to the outgoing null geodesic equation, which terminate in the past at the singularity and in future go to a distant observer in the spacetime, and if so under what conditions these exist. Let x_0 be the tangent to the null geodesics in (R, u) plane, at the central singularity, then it is given by,

$$x_0 = \lim_{t \rightarrow t_s} \lim_{r \rightarrow 0} \frac{R}{u} = \left. \frac{dR}{du} \right|_{t \rightarrow t_s; r \rightarrow 0} \quad (87)$$

Using Eq. (86), we get,

$$x_0^{\frac{3}{2}} = \frac{7}{4} \sqrt{M_0} \mathcal{X}_2(0) \quad (88)$$

In the (R, u) plane, the null geodesic equation will be,

$$R = x_0 u \quad (89)$$

while in the (t, r) plane, the null geodesic equation near the singularity will be,

$$t - t_s(0) = x_0 r^{\frac{7}{3}} \quad (90)$$

It follows that if $\mathcal{X}_2(0) > 0$, then that implies that $x_0 > 0$, and we then have radially outgoing null geodesics coming out from the singularity, making the *central singularity locally visible*. On the other hand, if $\mathcal{X}_2(0) < 0$, we will have a *black hole solution*. We have, however, already seen in Eq. (83), that the value of $\mathcal{X}_2(0)$ entirely depends upon the initial data and the cosmological term Λ . Given any Λ , the initial data can always be chosen such that the end state of the collapse would be either a naked singularity or a black hole. Hence, it follows that *for both positive and negative cosmological constants, a naked singularity can occur as the final end state of gravitational collapse*.

We noted here earlier that dust collapse models have been analyzed in the presence of a cosmological constant. The nature of the final singularity in that case has been analyzed using the so called ‘roots’ method (Deshingkar

S. S. *et al* [7]). We show below that the different treatment we have used above to deal with the collapsing clouds with pressure included, arrives at similar conclusions when specialized to the dust case.

The Einstein equations for *dust* may be obtained by putting $p_\theta = 0$ in the tangential pressure fluid model above. They take the form,

$$\rho(r, t) = \frac{\mathcal{M}'_d(r)}{R^2 R'} \quad (91)$$

$$\dot{R}^2 = \frac{\mathcal{M}_d(r)}{R} + f(r) + \frac{\Lambda R^2}{3} \quad (92)$$

where $f(r) = r^2 b_0(r)$ is the dust *energy free function*. Since we are interested in collapse we must have $\dot{R} \leq 0$. Then we have from the above equations,

$$\sqrt{R} \dot{R} = -\sqrt{\mathcal{M}_d(r) + f(r)R + \frac{\Lambda R^3}{3}} \quad (93)$$

Using the scaling freedom we may write,

$$\mathcal{M}_d(r) = \int_0^r \rho(0, r) r^2 dr = \rho^{avg}(r) r^3 \quad (94)$$

Rewriting equation (93) in terms of the scale factor v ,

$$\sqrt{v} \dot{v} = -\sqrt{\rho^{avg}(r) + \frac{f(r)v}{r^2} + \frac{\Lambda v^3}{3}} \quad (95)$$

Define $\frac{f(r)v}{r^2}$ as $L(r, v)$ which is at least a C^2 function of its arguments. Then $t_s(r, v)$ is readily calculated from (95) to be,

$$t_s(r, v) = \int_v^1 \frac{\sqrt{v} dv}{\sqrt{\rho^{avg}(r) + L(v, r) + \frac{\Lambda v^3}{3}}} \quad (96)$$

The shell focusing singularity $R = 0$ occurs first for the comoving coordinate $r = 0$. The time of its formation is,

$$t_s(0) = \int_0^1 \frac{\sqrt{v} dv}{\sqrt{\rho^{avg}(0) + L(v, 0) + \frac{\Lambda v^3}{3}}} \quad (97)$$

As we did for the fluid model, we Taylor expand $t_s(r)$ near $r = 0$ to get,

$$t_s(r) = t_s(0) + r \mathcal{X}_1(0) + r^2 \mathcal{X}_2(0) + \mathcal{O}(r^3) \quad (98)$$

where $\mathcal{X}_1(v) = 0$ (assuming smooth initial data), and

$$\mathcal{X}_2(v) = - \int_v^1 \frac{B_d''(0, v) \sqrt{v} dv}{B_d(0, v)^{3/2}} \quad (99)$$

with $B_d(r, v) = \rho^{avg}(r) + L(v, r) + \frac{\Lambda v^3}{3}$. Again, it needs to be analyzed whether there exists future directed null

geodesics that end at the singularity in the past. Consider the marginally bound case ($f(r) = 0$). Then the equation of the null geodesic is,

$$\frac{dt}{dr} = R' \quad (100)$$

In terms of (u, r) it is written as,

$$\frac{dR}{du} = \frac{R'}{\alpha r^{\alpha-1}} \left(1 - \sqrt{\frac{\mathcal{M}_d(r)}{R} + \frac{\Lambda R^2}{3}}\right) \quad (101)$$

Choosing $\alpha = \frac{7}{3}$,

$$\frac{dR}{du} = \frac{3}{7} \left(\frac{\sqrt{v} v'}{\sqrt{\frac{R}{u}}} + \frac{R}{u} \right) \left(1 - \sqrt{\frac{\mathcal{M}_d(r)}{R} + \frac{\Lambda R^2}{3}}\right) \quad (102)$$

Taking the limits as before, the final expression becomes,

$$\frac{4}{7} x_0^{\frac{3}{2}} = \sqrt{\rho_0^{avg}} \mathcal{X}_2(0) \quad (103)$$

It follows that for both positive and negative values of Λ , there exists initial data that may give $\mathcal{X}_2(0) > 0$. Thus the dust central singularity may be locally visible, even when there is a non-zero cosmological constant present. It may thus be claimed that the dust central naked singularity is not precluded by Λ .

VII. CONCLUSIONS

The final outcome of gravitational collapse is one of the most important open problems in gravitation theory, and the study of the fluid with vanishing radial pressure, and also the dust collapse models here indicate that the presence of a non-vanishing Λ cannot, in any conclusive manner, act as a cosmic censor. Although the value of the null geodesic tangents are modified by the presence of Λ , there are still cases where the singularity is locally visible depending on the initial data. The global visibility of such a singularity (null rays from the singularity reaching an asymptotic observer) which is locally naked, will depend on the overall behaviour of the various functions concerned which appear in the analysis. But as such general relativity has no natural length scale attached to

it, and so a locally visible singularity may violate the cosmic censorship as effectively as a globally visible singularity. Moreover, it is seen from various examples that if a singularity is locally naked, then one can always make it globally visible by suitable choice of the allowed functions, or by restricting properly the boundary of the cloud ([10]). We also note that studies pertaining to radiation collapse (of a *Type II* matter field [12]) in spacetimes with Λ , exist in literature [22]. Such studies also support the conclusion that both collapse endstates, namely a black hole or a naked singularity are possible in the presence of a non-zero Λ .

The currently observed accelerated expansion of the universe, and also the AdS/CFT conjecture in string theories have generated considerable interest in spacetime scenarios with a cosmological constant. We studied here dynamical collapse of a tangential pressure fluid model with non-vanishing Λ . Various aspects of gravitational collapse with Λ have been discussed in the context of the fluid model which include dynamics, junction conditions, apparent horizons, and nature of the singularity. The question of cosmic censorship in the collapse model is discussed, and it is also shown that when the pressures are put to zero in the model it reduces to the dust case.

The presence of a finite cosmological constant, it is seen, affects the dynamical collapse of the fluid in many ways. To summarize,

1. Λ plays a role in deciding the dynamical evolution of a shell. The conditions for singularity formation and bounce are derived.
2. The matching conditions for the interior and exterior spacetimes were discussed, assuming that the exterior is asymptotically AdS or dS.
3. The apparent horizon conditions and the time of horizon formation were studied in brief for $\Lambda \neq 0$.
4. The effect of Λ on the nature of the central singularity was analyzed. The condition for local visibility was derived and it was seen that even for non-vanishing Λ both black hole and naked singularity are possible end states of collapse.

Acknowledgments

A.M. would like to acknowledge support from TIFR, Mumbai during the Visiting Students Research Programme 2004 and thanks Ashutosh Mahajan for discussions.

-
- [1] Oppenheimer J. R. and Snyder H. , Phys. Rev, **56**, p.455 (1939).
 - [2] Penrose R., in *Black holes and relativistic stars*, ed. Wald R., University of Chicago Press (1998); Krolak A., Prog. Theor. Phys. Suppl., **136**, p. 45 (1999); Joshi P. S., Modern Physics Letters A, Vol.17, Nos. 15-17, 1067-1079 (2002) .
 - [3] Weinberg S., [astro-ph/9610044]; Carrol S. M., Living

- Rev. Rel. 4 (2001) 1 [astro-ph/0004075].
- [4] Perlmutter S. *et al.* (Supernova Cosmology Project Collaboration), Astrophys. Journ. **483**, p.565 (1997); Perlmutter S. *et al.* (Supernova Cosmology Project Collaboration), Nature **391**, p.51 (1998); Garnavich P. M. *et al.* (Hi-Z Supernova Team Collaboration), Astrophys. J. **493**, L53 (1998)
- [5] Maldacena J., Adv. Theor. Phys. **2** (1998) 231,

- [hep-th/9711200]; Aharony O., Gubser S., Maldacena J., Ooguri H., and Oz Y., Phys. Rept. 323 (2000) 183, [hep-th/9905111].
- [6] Krasinski A., *Inhomogeneous cosmological models*, Cambridge Univ. Press (1997)
- [7] Deshingkar S. S., Jhingan S., Chamorro A., and Joshi P. S., Phys.Rev. D **63** (2001) 124005; Cissoko M., Fabris J. C., Gariel J., Denmat G. L., and Santos N. O. , (1998) [gr-qc/9809057]; Markovic D. and Shapiro S. L., Phys. Rev D, **61** (2000); Lake K., Phys. Rev. D **62**, 027301 (2000).
- [8] Mahajan A., Goswami R. and Joshi P. S., Class. Quant. Grav. 22 (2005) 271-282.
- [9] Magli G., Class. Quant. Grav. **14** (1997) 1937; Class. Quant. Grav. **15** (1998) 3215; Goncalves S. M. C. V. , Jhingan S., Magli G., Phys. Rev. D **65** (2002) 064011; Harada T., Nakao K. and Eguchi H., Class.Quantum Grav. **16** (1999) 2785-2796; Goswami R. and Joshi P. S., Class. Quantum Grav.**19**, 5229 (2002).
- [10] Joshi P. S., *Global Aspects in Gravitation and Cosmology*, Oxford Univ. Press (1993).
- [11] Lemaitre G., Ann. Soc. Sci. Bruxelles I, **A53**,51(1933); H. Bondi, Mon. Not. Astron. Soc., **107**, 343 (1947); Tolman R. C., Proc. Nat. Acad. Sciences, **20**, p.169 (1934).
- [12] Hawking S. W. and Ellis G. F. R. , *The Large Scale Structure of Space-Time* (Cambridge, 1973).
- [13] Misner C. and Sharp D., Phys. Rev.**136**, B571 (1964).
- [14] Einstein A., Ann. Math. **40**, 4 922 (1939); Datta B. K., Gen. Relat. Grav. **1**, 19 (1970); Bondi H., Gen. Relat. Grav. **2**, 321 (1971).
- [15] Dickson L. E., *New First Course in the Theory of Equations*, John Wiley & sons, New York.
- [16] Darmon G., *Memorial des Sciences Mathematiques* (Paris,Gauthier-Villars)Fasc. 25 (1927).
- [17] Bonnor W. B. and Vickers P. A., Gen. Rel. Grav., **13**,29 (1981)
- [18] Israel W., Nuovo Cimento B, **44B**, 1 (1966).
- [19] Cahill M. E. and McVittie G. C., J.Math.Phys., **11**, 1382 (1970).
- [20] Shiromizu T., Nakao K., Kodama H., and Maeda K., Phys. Rev. D **47**,R3099 (1993); Hayward S. M., Shiromizu T. and Nakao K., Phys.Rev. D, **49**, 5080 (1994).
- [21] Goswami R. and Joshi P. S., Phys. Rev. D, **69**, (2004) 044002, [gr-qc/0212097]; Joshi P. S. and Dwivedi I. H., Phys. Rev. D, **47**, p.5357 (1993)
- [22] Lemos J. P. S., Phys.Rev.D **59** : 044020 (1999) [gr-qc/9812078]; Wagh S. M. and Maharaj S. D., Gen. Rel. Grav., Vol **31**, No 7 (1999).